

A PARAMETRIZATION OF THE GROUP OF SYMPLECTIC MATRICES AND ITS APPLICATIONS

Froilán M. Dopico

Departamento de Matemáticas
Universidad Carlos III de Madrid
Spain

Joint work with Charles R. Johnson, The College of William and Mary,
Williamsburg, VA, USA

Symplectic Matrices

$$J = \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix} \quad \text{where } I_n \text{ } n \times n \text{ identity matrix}$$

Definition: $S \in \mathbb{R}^{2n \times 2n}$ is symplectic if $S^T J S = J$

We will consider often the partition of symplectic S

$$S = \begin{bmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{bmatrix} \quad \text{with } S_{ij} \in \mathbb{R}^{n \times n}$$

The symplectic group is important in theory and applications (Hamiltonian mechanics, control,....)

The problem to be solved

Symplectic matrices are **implicitly defined** as solutions of the nonlinear matrix equation $S^T J S = J$

This makes difficult to work with them both in theory and in numerical algorithms.

OUR GOAL: To present an **explicit description (parametrization)** of the group of symplectic matrices, i.e., to find the set of solutions of

$$S^T J S = J \text{ where } S \text{ unknown}$$

and to apply this parametrization to different problems.

Outline of the talk

1. Previous results
2. Parametrization
3. Subparametrization problems
4. Description of Doubly structured sets (symplectic and other property):
 - LU factorizations of symplectic
 - Positive definite symplectic
 - Positive elementwise symplectic
 - TN, TP, oscillatory symplectic
 - Symplectic M-Matrices
5. Conclusions and Open problems

Previous I: A result by Mehrmann (SIMAX, 1988)

Theorem: Let $S = \begin{bmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{bmatrix}$ be symplectic with S_{11} non-singular. Then

$$S = \begin{bmatrix} I & 0 \\ S_{21}S_{11}^{-1} & I \end{bmatrix} \begin{bmatrix} S_{11} & 0 \\ 0 & S_{11}^{-T} \end{bmatrix} \begin{bmatrix} I & S_{11}^{-1}S_{12} \\ 0 & I \end{bmatrix}$$

where

1. The three factors are symplectic.
2. $S_{21}S_{11}^{-1}$ and $S_{11}^{-1}S_{12}$ are symmetric.

Let us combine this with three trivial facts...

Parametrization with nonsingular (1,1)-block

1. $\begin{bmatrix} I & 0 \\ X & I \end{bmatrix}$ is symplectic if and only if $X = X^T$.
2. $\begin{bmatrix} G & 0 \\ 0 & Y \end{bmatrix}$ is symplectic if and only if $Y = G^{-T}$.
3. Products and transposes of symplectic are symplectic.

Theorem: The set of symplectic matrices with nonsingular S_{11} is

$$\mathcal{S}^{(1,1)} = \left\{ \underbrace{\begin{bmatrix} G & GE \\ CG & G^{-T} + CGE \end{bmatrix}} : \begin{array}{l} G \text{ nonsingular} \\ C = C^T, E = E^T \end{array} \right\}$$
$$\begin{bmatrix} I & 0 \\ C & I \end{bmatrix} \begin{bmatrix} G & 0 \\ 0 & G^{-T} \end{bmatrix} \begin{bmatrix} I & E \\ 0 & I \end{bmatrix}$$

Parametrization with nonsingular (1,1)-block

Theorem: The set of symplectic matrices with nonsingular S_{11} is

$$\mathcal{S}^{(1,1)} = \left\{ \underbrace{\begin{bmatrix} G & GE \\ CG & G^{-T} + CGE \end{bmatrix}} : \begin{array}{l} G \text{ nonsingular} \\ C = C^T, E = E^T \end{array} \right\}$$
$$\begin{bmatrix} I & 0 \\ C & I \end{bmatrix} \begin{bmatrix} G & 0 \\ 0 & G^{-T} \end{bmatrix} \begin{bmatrix} I & E \\ 0 & I \end{bmatrix}$$

$2n^2 + n$ free parameters in this parametrization. This is precisely the dimension of the symplectic group.

What happens if S_{11} is singular?

The group of symplectic matrices

Theorem: The group of symplectic matrices is

$$S = \left\{ Q \begin{bmatrix} G & GE \\ CG & G^{-T} + CGE \end{bmatrix} : \right.$$

$$\left. \begin{array}{l} G \text{ nonsingular} \\ C = C^T, E = E^T \\ Q \text{ product of symplectic interchanges} \end{array} \right\}$$

REMARK: Given a symplectic matrix, Q may be not unique, then the previous description is not a *rigorous parametrization*. The nonuniqueness of Q can be useful for numerical purposes.

Subparametrization Problems (I)

1. Parametrization of symplectic matrices with $\text{rank}(S_{11}) = k$. This set depends on $2n^2 + n - \frac{(n-k)^2 + (n-k)}{2}$ parameters.

SIMILAR FOR ANY OTHER BLOCK

2. Any matrix can be S_{11} of a symplectic. If S_{11} is fixed and has $\text{rank}(S_{11}) = k$ the set of symplectic matrices compatible can be parametrized and depends on $\frac{n^2+n}{2} + \frac{k^2+k}{2} + n(n-k)$ parameters.

3. $\mathcal{S}^{(1,1)}$ is a dense subset of \mathcal{S} and sequences can be explicitly constructed.

Subparametrization Problems (II)

4. Parametrization of the set of $2n \times n$ matrices that can be $\begin{bmatrix} S_{11} \\ S_{21} \end{bmatrix}$ of a symplectic.

5. Parametrization of the set of symplectic matrices with given $\begin{bmatrix} S_{11} \\ S_{21} \end{bmatrix}$. This is an affine subspace in $\mathbb{R}^{2n \times 2n}$ of dimension $\frac{n^2+n}{2}$.

6. Any $A \in \mathbb{R}^{(n+1) \times (n+1)}$ can be $S(1:n+1, 1:n+1)$ of a symplectic S except by the fact that $a_{n+1, n+1}$ is determined by the other entries.

and more...

LU factorizations of Symplectic Matrices (I)

Theorem: Let $S = \begin{bmatrix} S_{11} & S_{12} \\ S_{21} & S_{22} \end{bmatrix}$ be symplectic. Then

1. S has LU factorization if and only if S_{11} and S_{11}^{-T} have LU factorizations.
2. S has LU factorization if and only if S_{11} is nonsingular and has LU and UL factorizations.
3. S has LU factorization if and only if $\det S_{11}(1 : k, 1 : k) \det S_{11}(k : n, k : n) \neq 0 \quad k = 1 : n$

to be continued....

LU factorizations of Symplectic Matrices (II)


4. If $S_{11} = L_{11}U_{11}$ and $S_{11}^{-T} = L_{22}U_{22}$ are LU factorizations, then the LU factorization of S is

$$S = \begin{bmatrix} L_{11} & 0 \\ S_{21}U_{11}^{-1} & L_{22} \end{bmatrix} \begin{bmatrix} U_{11} & L_{11}^{-1}S_{12} \\ 0 & U_{22} \end{bmatrix}$$

5. The LU factors of S are symplectic if and only if S_{11} is diagonal.

Symplectic LU-like factorization

$$S = \begin{bmatrix} L_{11} & 0 \\ S_{21}U_{11}^{-1} & L_{11}^{-T} \end{bmatrix} \begin{bmatrix} U_{11} & L_{11}^{-1}S_{12} \\ 0 & U_{11}^{-T} \end{bmatrix}$$


upper triang. lower triang.

Symplectic Positive Definite (PD) Matrices

Theorem: Let $S = \begin{bmatrix} S_{11} & S_{21}^T \\ S_{21} & S_{22} \end{bmatrix}$ be symmetric and symplectic. Then

1. S is PD if and only if S_{11} is PD.

2. The set of PD symplectic matrices is

$$\mathcal{S}^{\text{PD}} = \left\{ \begin{bmatrix} G & GC \\ CG & G^{-1} + CGC \end{bmatrix} : \begin{array}{l} G \text{ positive definite} \\ C = C^T \end{array} \right\}$$

\mathcal{S}^{PD} depends on $n^2 + n$ parameters.

3. Every PD symplectic matrix $S = HH^T$ with H symplectic.

Symplectic Matrices with positive entries

Set of symplectic matrices with nonsingular S_{11}

$$\mathcal{S}^{(1,1)} = \left\{ \left[\begin{array}{cc} G & GE \\ CG & G^{-T} + CGE \end{array} \right] : \begin{array}{l} G \text{ nonsingular} \\ C = C^T, E = E^T \end{array} \right\}$$

This allows us to generate symplectic matrices with positive entries (contrast with orthogonal matrices).

START by choosing arbitrary $G > 0$, $C > 0$, and $\tilde{E} > 0$ with positive entries. So $CG\tilde{E} > 0$.

THEN, a number $\alpha > 0$ is chosen such that

$$\alpha CG\tilde{E} + G^{-T} > 0.$$

FINALLY: $E \equiv \alpha\tilde{E}$

Totally Nonnegative (TN) Symplectic Matrices (I)

DEFINITIONS:

Matrices with all minors nonnegative (**positive**) are called TN (**totally positive TP**) matrices.

If A is TN and A^k TP for some positive integer k then A is called **OSCILLATORY**.

Applications in mechanical oscillatory problems

Totally Nonnegative (TN) Symplectic Matrices (II)

I is symplectic and TN.

Are there symplectic TP matrices?

Are there symplectic oscillatory matrices?

What is the set of symplectic TN matrices?

Theorem (the 2×2 case): $S \in \mathbb{R}^{2 \times 2}$ is symplectic and TP if and only if $\det S = 1$ and $s_{ij} > 0$ for all (i, j) .

If any three positive entries such that $s_{11}s_{22} > 1$ are chosen then the remaining entry is obtained from $\det S = 1$.

Totally Nonnegative (TN) Symplectic Matrices (III)

Theorem: Let $S \in \mathbb{R}^{2n \times 2n}$ with $n > 1$ be symplectic. Then

1. S is not TP.
2. S is not oscillatory.

Sketch of the Proof: LU factorization of S is

$$S = \begin{bmatrix} L_{11} & 0 \\ S_{21}U_{11}^{-1} & L_{22} \end{bmatrix} \begin{bmatrix} U_{11} & L_{11}^{-1}S_{12} \\ 0 & U_{22} \end{bmatrix} \quad \text{where} \quad \begin{array}{l} S_{11} = L_{11}U_{11} \\ S_{11}^{-T} = L_{22}U_{22} \end{array}$$

If S TP then S_{11} is TP and the LU factors of S are triangular TP. Therefore L_{22} and U_{22} are triangular TP and S_{11}^{-T} is TP.

CONTRADICTION!!, S_{11} TP implies that S_{11}^{-T} has negative entries.

Totally Nonnegative (TN) Symplectic Matrices (IV)

Theorem: The set of $2n \times 2n$ ($n > 1$) symplectic and TN matrices is

$$\mathcal{S}^{\text{TN}} = \left\{ \begin{bmatrix} D & 0 \\ 0 & D^{-1} \end{bmatrix} : D = \begin{bmatrix} \lambda_1 & & \\ & \cdots & \\ & & \lambda_n \end{bmatrix} \lambda_i > 0 \right\}$$

Symplectic M-Matrices (I)

Definition: $A \in \mathbb{R}^{n \times n}$ is a **M-Matrix** if $a_{ij} \leq 0$ for $i \neq j$ and $\operatorname{Re}(\lambda) > 0$ for every eigenvalue λ of A .

Theorem: The set of $2n \times 2n$ symplectic M-Matrices is

$$\mathcal{S}^M = \left\{ \left[\begin{array}{cc} D & DK \\ HD & D^{-1} + HDK \end{array} \right] : \left. \begin{array}{l} D \text{ positive diagonal} \\ H = H^T \leq 0 \\ K = K^T \leq 0 \\ HDK \text{ diagonal} \end{array} \right\}$$

Symplectic M-Matrices (II)

Given an arbitrary $H = H^T \leq 0$, the matrices $K = K^T \leq 0$ such that HDK is *diagonal* can be easily determined. For instance if $h_{12} = h_{21} \neq 0$:

$$H = \begin{bmatrix} & x \\ x & \end{bmatrix} \longrightarrow K = \begin{bmatrix} 0 & ? & 0 & 0 & 0 \\ ? & 0 & 0 & 0 & 0 \\ 0 & 0 & & & \\ 0 & 0 & & & \\ 0 & 0 & & & \end{bmatrix}$$

The ? that remain in K after this process is repeated for all entries $h_{ij} = h_{ji} \neq 0$, are free parameters in $K = K^T \leq 0$ for a given H .

Conclusions and Open Problems

- An explicit description of the group of symplectic matrices has been introduced.
- It allows to solve very easily many theoretical questions.
- Perturbation theory with respect the symplectic parameters? Interesting properties?
- How to compute the parametrization in a stable an efficient way if we are given the entries of a symplectic matrix?
- Is it possible to get a rank revealing factorization?
- Have these symplectic parameters an intrinsic meaning?